


Iowa County Health Snapshot Series

99 county reports prepared for the Iowa State University Extension and Outreach Community and Economic Development Unit



May 2024

By Matthew Voss & Shawn Dorius

This project was supported by funding from Iowa State University's Extension and Outreach Community and Economic Development Unit via an award from Iowa State University's Presidential Strategic Initiative Plan 2022-2031.

Iowa County Health Snapshot Series

Table of contents

| Health in Marion County: Overvie Strengths | | | | | | | | | | | | | | | | | | 3 |
|--|------|----|----|-------|---|-------|---|---|-------|-------|---|-------|---|---|---|-------|---|---------|
| Opportunities | | | | | | | | | | | | | | | | | | |
| • • | | | | | | | | | | | | | | | | | | |
| How to Use this Report | | | | | | | | | | | | | | | | | | |
| Using Story County as an E | :xai | пр | ie | • | • | • | • | • | • | • | • | • | • | • | • | • | • | 6 |
| Health Status | | | | | | | | | | | | | | | | | | 7 |
| Poor General Health | | | | | | | | | | | | | | | | | | . 7 |
| Poor Physical Health | | | | | | | | | | | | | | | | | | . 8 |
| Poor Mental Health | | | | | | | | | | | | | | | | | | |
| Health Risk Behaviors | | | | | | | | | | | | | | | | | | 9 |
| Binge Drinking | | | | | | | | | | | | | | | | | | g |
| Current Smoking | | | | | | | | | | | | | | | | | | |
| Physical Inactivity | | | | | | | | | | | | | | | | | | |
| Sleep <7 hours | | | | | | | | | | | | | | | | | | |
| Health Outcomes | | | | | | | | | | | | | | | | | | 12 |
| Arthritis | | | | | | | | | | | | | | | | | | |
| High Blood Pressure | | | | | | | | | | | | | | | | | | |
| Cancer (except skin) | | | | | | | | | | | | | | | | | | _ |
| Current Asthma | | | | | | | | | | | | | | | | | | |
| Coronary Heart Disease . | | | | | | | | | | | | | | | | | | |
| COPD | | | | | | | | | | | | | | | | | | |
| Depression | | | | | | | | | | | | | | | | | | |
| Diabetes | | | | | | | | | | | | | | | | | | |
| High Cholesterol | | | | | | | | | | | | | | | | | | _ |
| Chronic Kidney Disease . | | | | | | | | | | | | | | | | | | |
| Obesity | | | | | | | | | | | | | | | | | | |
| Stroke | | | | | | | | | | | | | | | | | | |
| All Teeth Lost | | | | | | | | | | | | | | | | | | |
| All feetil Lost | | • | • | • | • | • | • | • | • | • | • | • | • | • | • | • | • | 10 |
| Prevention | | | | | | | | | | | | | | | | | | 19 |
| Lack of Health Insurance . | | | | | | | | | | | | | | | | | | 19 |
| Taking BP Medication | | | | | | | | | | | | | | | | | | 20 |
| Cervical Cancer Screening | | | | | | | | | | | | | | | | | | 20 |
| Annual Checkup | | | | | | | | | | | | | | | | | | 21 |
| Cholesterol Screening | | | | | | | | | | | | | | | | | | |

| | Colorectal Cancer Screening | 22 24 24 |
|----------|---|----------------------------|
| Disabili | Hearing disability Vision disability Cognitive disability Mobility disability Self-care disability Independent living disability Any disability | 28 28 29 29 30 |
| Append | dix 1: CDC PLACES Data | 31 |
| Append | dix 2: Marion County PLACES Data | 32 |
| Append | dix 3: Micropolitan Counties | 34 |
| Append | dix 4: County Maps | 35 |

Health in Marion County: Overview

Dear Community Members & Stakeholders of Marion County,

We are pleased to share the Iowa County Health Snapshot series, a set of reports designed in partner-ship with Iowa State University's Community Economic Development to assess the overall landscape of health and well-being across Iowa's 99 counties. The following pages include the health snapshot for your county. The snapshot can be used to see how Marion County ranks across various health measures compared to the rest of the state and similarly-sized counties. This comparison allows us to identify Marion County's strengths and opportunities for growth by leveraging data from the Centers for Disease Control and Prevention PLACES dataset.

The report includes 37 health measures organized topically by general health status, risky behavior, health outcomes, prevention, and disability. We provide a general overview of each health topic, followed by a description of each health measure. Health measure descriptions include crude and age-adjusted rates and a ranking that tells you how your county ranks compared to all lowa counties and counties of similar size to Marion County. Crude rates tell you the raw, un-adjusted percent of the population experiencing the health measure. Age-adjusted rates account for the age distribution present in your county and presents what the rate would be if your county had a standard age distribution. Crude rates can provide a general comparison between counties and age-adjusted rates can give a fairer comparison between counties with different age structures.

Strengths

Take a look at the top strengths of your county! Below are unadjusted and age-adjusted health measures that represent your county's strengths. We recommend using continued investment in these areas to maintain them as strengths, as well as building on them to improve in other health domains.

| | Crude | | Age | -Adjusted | |
|----------------------|--------------------------|-----------------------------|------------------------|--------------------------|-----------------------------|
| Health Domain | Statewide County Rank | Micropolitan County Rank | Health Domain | Statewide County Rank | Micropolitan County Rank |
| Dental Visit | 2nd | 1st | Dental Visit | 2nd | 1st |
| Mammography | 4th | 1st | Stroke | 3rd | 1st |
| Current Asthma | 4th | 2nd | Chronic Kidney Disease | 3rd | 1st |
| Self-care Disability | 7th | 1st | Mammography | 4th | 1st |
| Vision Disability | 8th | 1st | Current Asthma | 4th | 2nd |

Opportunities

Below are health measures that represent potential areas for growth in your county. We recommend increasing investment in these areas to support community health and well-being.

| | Crude | | Age | -Adjusted | |
|-----------------------|--------------------------|-----------------------------|--|--------------------------|-----------------------------|
| Health Domain | Statewide County Rank | Micropolitan County Rank | Health Domain | Statewide County Rank | Micropolitan County Rank |
| Binge Drinking | 87th | 15th | High Cholesterol | 91st | 11th |
| Taking BP Medication | 84th | 13th | Binge Drinking | 90th | 14th |
| Depression | 62nd | 6th | Taking BP Medication | 82nd | 15th |
| Cholesterol Screening | 60th | 7th | Depression | 48th | 6th |
| High Cholesterol | 39th | 6th | Core preventive services for older men | 31st | 6th |

Please review the rest of the report to learn more about your county and reach out to us with questions about the report or for recommendations to improve the overall health and well-being in Marion County.

Sincerely,

Shawn F. Dorius, PhD Iowa State University

Show Di

Public Science Collaborative



IOWA STATE UNIVERSITY
Extension and Outreach
Community and Economic Development

This project was supported by funding from Iowa State University's Extension and Outreach Community and Economic Development Unit, via an award from Iowa State University's Presidential Strategic Initiative Plan 2022-2031.

How to Use this Report


The purpose of the Iowa County Health Snapshot Series is to provide an overview of health in Marion County that can be used to help inform health-related interventions and programming. This report contains five sections, including the overall health status of Marion County, a review of risky behaviors in the county, and information on various health outcomes, prevention measures, and disability needs. Each section includes an overview of the health domain, a breakdown of its underlying health measures, and data to see how Marion County compares to the rest of the state and similarly-sized counties.

With a population of 33,380, Marion County's similarly sized counties are defined as micropolitan counties. In lowa, micropolitan counties have an average population of 30,839, and there are 15 counties that are considered micropolitan counties. To learn more about the county population classifications used in this report, please see the information below the table as well as Appendix 3.

| Group | Population Avg. | Number of Counties |
|--------------------------|-----------------|--------------------|
| Metropolitan Core | 175,797 | 9 |
| Micropolitan | 30,839 | 15 |
| Metropolitan Non-Core | 29,821 | 12 |
| Rural | 12,547 | 63 |
| Total | | 99 |


The counties included in each category above are determined by the United States Office of Management and Budget. Iowa has counties in nine metropolitan statistical areas and 15 micropolitan statistical areas. The counties identified as metropolitan and micropolitan can be found in the link and the maps below. Metropolitan core counties are those counties in metropolitan areas that have the city that defines the metropolitan area (e.g., Des Moines and Polk County for the Des Moines Metropolitan Area). Metropolitan non-core counties are those that fall within the metropolitan area, but do not have the core city of the metropolitan area (e.g., Dallas and Jasper Counties in the Des Moines Metropolitan Area).

https://www.icip.iastate.edu/maps/refmaps/cofips



Using Story County as an Example

The graphic below is an example of the details included for each health measure. The table in the upper right shows your county's crude and age-adjusted rate estimates alongside their averages across all lowa counties and similarly-sized counties. The images below show the distribution of the health measure across counties. The vertical lines correspond with the values in the table to show the estimate for the specific county compared to counties across the state. The solid black line in each plot represents the county rate, with the dashed lines indicating the average rate for all lowa counties and the dotted lines showing the average rate for similarly-sized counties.



The example graphic above displays the poor mental health rate in Story County, Iowa. The left plot shows the crude rate of poor mental health in the county. Crude rates are the raw, un-adjusted percent of the population experiencing the health measure. Notice how the solid black line is closer to the right of the plot than the dashed and dotted lines; this indicates that Story County has a higher crude rate of poor mental health than most counties in the state. However, the age-adjusted rate plot on the right shows the solid line closer to the peak of the distribution and averages for the state and similarly-sized counties, indicating that when accounting for Story County's unique age distribution, Story County's poor mental health rate is not a significant outlier, and is estimated to be lower than the statewide average. Accounting for crude and age-adjusted rates is critical to help inform health interventions as age-adjusted rates can help contextualize the health measure within the population.

Health Status


This section reflects measures for overall health, including general, physical, and mental health. These measures are a valuable way to see a simple summary of the health of a country's population. The questions to calculate these measures ask respondents to evaluate their health status. General health is evaluated on a 5-point scale where respondents can report their health status as one of "excellent," "very good," "good," "fair," or "poor," from which the respondents who answer "fair" or "poor" are included in the measure. Physical and mental health are evaluated as the rate of respondents who say they have had 14 or more days in the last 30 days in which their physical or mental health was not good. The URL below includes more information about each variable.

https://www.cdc.gov/places/measure-definitions/health-status/index.html

Poor General Health

Marion County has a crude rate of 12.5% for fair or poor self-rated health status among adults aged >= 18 years, which ranks 11th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 11.5%, which ranks 9th out of 99 lowa counties (2nd out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 12.5% | 15.0% | 15.3% |
| Age-Adjusted | 11.5% | 13.3% | 14.1% |



Poor Physical Health

Marion County has a crude rate of 9.7% for physical health not good for >=14 days among adults aged >=18 years, which ranks 10th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 9.0%, which ranks 12th out of 99 lowa counties (2nd out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 9.7% | 11.1% | 11.1% |
| Age-Adjusted | 9.0% | 9.9% | 10.3% |



Poor Mental Health

Marion County has a crude rate of 13.9% for mental health not good for >=14 days among adults aged >=18 years, which ranks 35th out of 99 lowa counties (6th out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 14.8%, which ranks 15th out of 99 lowa counties (4th out of 15 similarly-sized counties).

| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 13.9% | 14.1% | 14.5% |
| Age-Adjusted | 14.8% | 15.5% | 15.5% |



Health Risk Behaviors


Health risk measures are measures of individual behaviors that can lead to poor health outcomes. These include binge drinking, current smoking, physical inactivity, and getting less than seven hours of nightly sleep. Risk behaviors are the most critical intervention points as they can help save lives and drastically improve health outcomes later in life. Each measure is calculated as the rate of respondents participating in a high level of risk behavior. The exact cutoff points for the behavior depend on the health area. The link below includes more information about each variable.

https://www.cdc.gov/places/measure-definitions/unhealthy-behaviors/index.html

Binge Drinking

Marion County has a crude rate of 20.3% for binge drinking among adults aged >=18 years, which ranks 87th out of 99 lowa counties (15th out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 22.5%, which ranks 90th out of 99 lowa counties (14th out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 20.3% | 18.5% | 18.1% |
| Age-Adjusted | 22.5% | 21.3% | 20.8% |



Current Smoking

Marion County has a crude rate of 14.7% for current smoking among adults aged >=18 years, which ranks 10th out of 99 lowa counties (2nd out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 15.7%, which ranks 14th out of 99 lowa counties (2nd out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 14.7% | 16.7% | 16.6% |
| Age-Adjusted | 15.7% | 17.8% | 17.5% |



Physical Inactivity

Marion County has a crude rate of 23.6% for no leisuretime physical activity among adults aged >=18 years, which ranks 11th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 22.6%, which ranks 14th out of 99 lowa counties (2nd out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 23.6% | 26.8% | 27.4% |
| Age-Adjusted | 22.6% | 25.1% | 25.7% |



Sleep <7 hours

Marion County has a crude rate of 29.8% for sleeping less than 7 hours among adults aged >=18 years, which ranks 38th out of 99 lowa counties (5th out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 30.9%, which ranks 31st out of 99 lowa counties (4th out of 15 similarly-sized counties).

| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 29.8% | 30.0% | 30.5% |
| Age-Adjusted | 30.9% | 31.4% | 31.6% |



Health Outcomes


This section includes measures about specific health outcomes such as chronic illness, mental health conditions, and risk factors for future disease. Health outcome measures include but are not limited to arthritis, high blood pressure, cancer, depression, and obesity. Each measure is the percentage of respondents reporting receiving the diagnosis by a doctor, nurse, or other health professional. The link below includes more information about each variable.

https://www.cdc.gov/places/measure-definitions/health-outcomes/index.html

Arthritis

Marion County has a crude rate of 25.2% for arthritis among adults aged >=18 years, which ranks 13th out of 99 lowa counties (3rd out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 21.4%, which ranks 6th out of 99 lowa counties (3rd out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 25.2% | 28.8% | 29.0% |
| Age-Adjusted | 21.4% | 23.2% | 23.8% |



High Blood Pressure

Marion County has a crude rate of 31.3% for high blood pressure among adults aged >=18 years, which ranks 16th out of 99 lowa counties (3rd out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 27.2%, which ranks 29th out of 99 lowa counties (4th out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 31.3% | 33.7% | 33.3% |
| Age-Adjusted | 27.2% | 27.9% | 27.7% |



Cancer (except skin)

Marion County has a crude rate of 7.8% for cancer (excluding skin cancer) among adults aged >=18 years, which ranks 18th out of 99 lowa counties (7th out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 6.4%, which ranks 26th out of 99 lowa counties (7th out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 7.8% | 8.4% | 8.1% |
| Age-Adjusted | 6.4% | 6.4% | 6.4% |



Current Asthma

Marion County has a crude rate of 9.0% for current asthma among adults aged >=18 years, which ranks 4th out of 99 lowa counties (2nd out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 9.1%, which ranks 4th out of 99 lowa counties (2nd out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 9.0% | 9.3% | 9.3% |
| Age-Adjusted | 9.1% | 9.5% | 9.5% |



Coronary Heart Disease

Marion County has a crude rate of 6.1% for coronary heart disease among adults aged >=18 years, which ranks 14th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 4.8%, which ranks 9th out of 99 lowa counties (1st out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 6.1% | 7.1% | 7.0% |
| Age-Adjusted | 4.8% | 5.2% | 5.2% |



COPD

Marion County has a crude rate of 6.5% for chronic obstructive pulmonary disease among adults aged >=18 years, which ranks 9th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 5.5%, which ranks 8th out of 99 lowa counties (1st out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 6.5% | 7.9% | 7.7% |
| Age-Adjusted | 5.5% | 6.4% | 6.3% |



Depression

Marion County has a crude rate of 17.9% for depression among adults aged >=18 years, which ranks 62nd out of 99 lowa counties (6th out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 18.7%, which ranks 48th out of 99 lowa counties (6th out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 17.9% | 17.6% | 18.0% |
| Age-Adjusted | 18.7% | 18.7% | 18.8% |



Diabetes

Marion County has a crude rate of 9.7% for diagnosed diabetes among adults aged >=18 years, which ranks 14th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 8.0%, which ranks 13th out of 99 lowa counties (2nd out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 9.7% | 11.0% | 11.0% |
| Age-Adjusted | 8.0% | 8.7% | 9.1% |



High Cholesterol

Marion County has a crude rate of 37.2% for high cholesterol among adults aged >=18 years who have been screened in the past 5 years, which ranks 39th out of 99 lowa counties (6th out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 30.9%, which ranks 91st out of 99 lowa counties (11th out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 37.2% | 37.5% | 37.4% |
| Age-Adjusted | 30.9% | 29.8% | 30.0% |



Chronic Kidney Disease

Marion County has a crude rate of 3.0% for chronic kidney disease among adults aged >=18 years, which ranks 9th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 2.5%, which ranks 3rd out of 99 lowa counties (1st out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 3.0% | 3.4% | 3.4% |
| Age-Adjusted | 2.5% | 2.7% | 2.7% |



Obesity

Marion County has a crude rate of 36.3% for obesity among adults aged >=18 years, which ranks 25th out of 99 lowa counties (5th out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 36.8%, which ranks 24th out of 99 lowa counties (5th out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 36.3% | 37.6% | 37.1% |
| Age-Adjusted | 36.8% | 37.9% | 37.4% |



Stroke

Marion County has a crude rate of 3.0% for stroke among adults aged >=18 years, which ranks 10th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 2.4%, which ranks 3rd out of 99 lowa counties (1st out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 3.0% | 3.5% | 3.5% |
| Age-Adjusted | 2.4% | 2.7% | 2.8% |



All Teeth Lost

Marion County has a crude rate of 8.5% for all teeth lost among adults aged >=65 years, which ranks 9th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 9.0%, which ranks 8th out of 99 lowa counties (1st out of 15 similarly-sized counties).

| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 8.5% | 10.4% | 11.1% |
| Age-Adjusted | 9.0% | 11.3% | 11.9% |



Prevention


Preventive measures are another practical intervention point for public health professionals as they allow for the early detection of health conditions and improve overall treatment efficacy. This section includes a review of your county's prevention measures, including preventive screenings and doctor visits. Besides the measure for lack of health insurance, all measures in this section are positive, where a higher rate is a strength for your county. The link below includes more information about each variable.

https://www.cdc.gov/places/measure-definitions/prevention/index.html

Lack of Health Insurance

Marion County has a crude rate of 6.6% for current lack of health insurance among adults aged 18–64 years, which ranks 9th out of 99 lowa counties (2nd out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 6.9%, which ranks 8th out of 99 lowa counties (1st out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 6.6% | 7.8% | 8.0% |
| Age-Adjusted | 6.9% | 8.3% | 8.5% |



Taking BP Medication

Marion County has a crude rate of 79.4% for taking medicine for high blood pressure control among adults aged >=18 years with high blood pressure, which ranks 84th out of 99 lowa counties (13th out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 58.6%, which ranks 82nd out of 99 lowa counties (15th out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 79.4% | 80.8% | 80.1% |
| Age-Adjusted | 58.6% | 59.4% | 59.9% |



Cervical Cancer Screening

Marion County has a crude rate of 83.8% for cervical cancer screening among adult women aged 21–65 years, which ranks 11th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 84.6%, which ranks 6th out of 99 lowa counties (1st out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 83.8% | 82.3% | 81.9% |
| Age-Adjusted | 84.6% | 83.0% | 82.5% |



Annual Checkup

Marion County has a crude rate of 76.5% for visits to doctor for routine checkup within the past year among adults aged >=18 years, which ranks 27th out of 99 lowa counties (5th out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 74.2%, which ranks 5th out of 99 lowa counties (2nd out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 76.5% | 75.9% | 75.4% |
| Age-Adjusted | 74.2% | 72.6% | 72.6% |



Cholesterol Screening

Marion County has a crude rate of 82.2% for cholesterol screening among adults aged >=18 years, which ranks 60th out of 99 lowa counties (7th out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 80.1%, which ranks 16th out of 99 lowa counties (1st out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 82.2% | 82.5% | 82.2% |
| Age-Adjusted | 80.1% | 78.9% | 79.1% |



Colorectal Cancer Screening

Marion County has a crude rate of 74.7% for colorectal cancer screening among adults aged 50–75 years, which ranks 20th out of 99 lowa counties (2nd out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 72.6%, which ranks 12th out of 99 lowa counties (1st out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 74.7% | 73.2% | 72.4% |
| Age-Adjusted | 72.6% | 70.3% | 69.5% |



Core preventive services for older men

Marion County has a crude rate of 44.5% for older adult men aged >=65 years who are up to date on a core set of clinical preventive services: flu shot past year, ppv shot ever, colorectal cancer screening, which ranks 30th out of 99 lowa counties (6th out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 44.9%, which ranks 31st out of 99 lowa counties (6th out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 44.5% | 42.7% | 43.6% |
| Age-Adjusted | 44.9% | 43.3% | 44.4% |



Core preventive services for older women

Marion County has a crude rate of 42.2% for older adult women aged >=65 years who are up to date on a core set of clinical preventive services: flu shot past year, ppv shot ever, colorectal cancer screening, and mammogram past 2 years, which ranks 14th out of 99 lowa counties (3rd out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 42.8%, which ranks 15th out of 99 lowa counties (3rd out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 42.2% | 38.4% | 39.2% |
| Age-Adjusted | 42.8% | 39.7% | 39.5% |



Dental Visit

Marion County has a crude rate of 74.3% for visits to dentist or dental clinic among adults aged >=18 years, which ranks 2nd out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 73.6%, which ranks 2nd out of 99 lowa counties (1st out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 74.3% | 66.0% | 64.2% |
| Age-Adjusted | 73.6% | 65.3% | 63.8% |



Mammography

Marion County has a crude rate of 77.5% for mammography use among women aged 50–74 years, which ranks 4th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 77.3%, which ranks 4th out of 99 lowa counties (1st out of 15 similarly-sized counties).

| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 77.5% | 73.9% | 73.4% |
| Age-Adjusted | 77.3% | 73.7% | 73.5% |



Disability


This section includes measures of physical and cognitive disabilities. Respondents were asked to report on six disabilities: hearing, vision, cognition, mobility, self-care, and independent living. The final measure reports the rate of respondents that answered yes to any of the previous six disabilities. The link below includes more information about each variable.

https://www.cdc.gov/places/measure-definitions/unhealthy-behaviors/index.html

Hearing disability

Marion County has a crude rate of 6.7% for hearing disability among adults aged >=18 years, which ranks 11th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 5.6%, which ranks 8th out of 99 lowa counties (1st out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 6.7% | 7.9% | 7.8% |
| Age-Adjusted | 5.6% | 6.1% | 6.2% |



Vision disability

Marion County has a crude rate of 3.5% for vision disability among adults aged >=18 years, which ranks 8th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 3.2%, which ranks 6th out of 99 lowa counties (1st out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 3.5% | 4.4% | 4.6% |
| Age-Adjusted | 3.2% | 3.9% | 4.2% |



Cognitive disability

Marion County has a crude rate of 10.3% for cognitive disability among adults aged >=18 years, which ranks 11th out of 99 lowa counties (2nd out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 10.6%, which ranks 11th out of 99 lowa counties (2nd out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 10.3% | 11.2% | 11.7% |
| Age-Adjusted | 10.6% | 11.9% | 12.1% |



Mobility disability

Marion County has a crude rate of 10.4% for mobility disability among adults aged >=18 years, which ranks 10th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 8.7%, which ranks 9th out of 99 lowa counties (1st out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 10.4% | 12.5% | 12.2% |
| Age-Adjusted | 8.7% | 9.9% | 10.4% |



Self-care disability

Marion County has a crude rate of 2.9% for self-care disability among adults aged >=18 years, which ranks 7th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 2.6%, which ranks 7th out of 99 lowa counties (1st out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 2.9% | 3.6% | 3.7% |
| Age-Adjusted | 2.6% | 3.1% | 3.3% |



Independent living disability

Marion County has a crude rate of 5.8% for independent living disability among adults aged >=18 years, which ranks 9th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 5.5%, which ranks 8th out of 99 lowa counties (1st out of 15 similarly-sized counties).


| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 5.8% | 6.9% | 7.2% |
| Age-Adjusted | 5.5% | 6.4% | 6.9% |



Any disability

Marion County has a crude rate of 23.3% for any disability among adults aged >=18 years, which ranks 11th out of 99 lowa counties (1st out of 15 similarly-sized counties). The age-adjusted rate for Marion County is 21.5%, which ranks 9th out of 99 lowa counties (1st out of 15 similarly-sized counties).

| Measure Type | Marion County | All Counties | Similarly- Sized Counties |
|--------------|------------------|-----------------|---------------------------------|
| Crude | 23.3% | 26.6% | 26.6% |
| Age-Adjusted | 21.5% | 23.6% | 24.4% |



Appendix 1: CDC PLACES Data

This report uses a novel dataset from the Centers for Disease Control and Prevention (CDC) called PLACES: Local Data for Better Health. The PLACES dataset uses small area estimation methods to provide estimates of health measures for counties, places, census tracts, and zip code tabulation areas. The estimates are created with a multilevel statistical modeling framework that utilizes CDC's Behavioral Risk Factor Surveillance System (BRFSS), Census population counts, and American Community Survey demographic and socioeconomic data.

Relevant links for CDC PLACES can be found below:

- PLACES Homepage
- Measure Definitions
- Methodology



Appendix 2: Marion County PLACES Data

| Category | Health Domain | Crude Rate Rank | Crude Rate (95% CI) | Age-Adj. Rate Rank | Age-Adj. Rate (95% CI) |
|--------------------------|--|--------------------|-----------------------|-----------------------|------------------------|
| Health Status | General Health | 11 | 12.5% (10.2% - 15.1%) | 9 | 11.5% (9.4% - 13.9%) |
| Health Status | Physical Health | 10 | 9.7% (8.0% - 11.4%) | 12 | 9.0% (7.4% - 10.6%) |
| Health Status | Mental Health | 35 | 13.9% (11.7% - 16.3%) | 15 | 14.8% (12.4% - 17.3%) |
| Health Risk Behaviors | Physical Inactivity | 11 | 23.6% (18.9% - 28.8%) | 14 | 22.6% (18.1% - 27.6%) |
| Health Risk Behaviors | Sleep <7 hours | 38 | 29.8% (28.1% - 31.4%) | 31 | 30.9% (29.0% - 32.6%) |
| Health Risk Behaviors | Current Smoking | 10 | 14.7% (11.3% - 18.5%) | 14 | 15.7% (11.8% - 19.8%) |
| Health Risk Behaviors | Binge Drinking | 87 | 20.3% (16.6% - 24.4%) | 90 | 22.5% (18.5% - 27.0%) |
| Health Outcomes | Stroke | 10 | 3.0% (2.5% - 3.5%) | 3 | 2.4% (2.1% - 2.8%) |
| Health Outcomes | Cancer (except skin) | 18 | 7.8% (7.0% - 8.8%) | 26 | 6.4% (5.7% - 7.2%) |
| Health Outcomes | Coronary Heart Disease | 14 | 6.1% (5.1% - 7.1%) | 9 | 4.8% (4.1% - 5.7%) |
| Health Outcomes | COPD | 9 | 6.5% (5.1% - 8.0%) | 8 | 5.5% (4.4% - 6.8%) |
| Health Outcomes | Arthritis | 13 | 25.2% (21.0% - 29.5%) | 6 | 21.4% (17.6% - 25.4%) |
| Health Outcomes | High Blood Pressure | 16 | 31.3% (26.7% - 35.8%) | 29 | 27.2% (23.0% - 31.5%) |
| Health Outcomes | Obesity | 25 | 36.3% (28.9% - 44.1%) | 24 | 36.8% (29.4% - 44.6%) |
| Health Outcomes | Diabetes | 14 | 9.7% (7.9% - 11.6%) | 13 | 8.0% (6.5% - 9.6%) |
| Health Outcomes | All Teeth Lost | 9 | 8.5% (5.2% - 12.3%) | 8 | 9.0% (5.6% - 13.0%) |
| Health Outcomes | Depression | 62 | 17.9% (14.3% - 21.9%) | 48 | 18.7% (14.9% - 22.8%) |
| Health Outcomes | High Cholesterol | 39 | 37.2% (32.4% - 42.3%) | 91 | 30.9% (26.1% - 36.0%) |
| Health Outcomes | Current Asthma | 4 | 9.0% (7.7% - 10.4%) | 4 | 9.1% (7.8% - 10.6%) |
| Health Outcomes | Chronic Kidney Disease | 9 | 3.0% (2.6% - 3.4%) | 3 | 2.5% (2.2% - 2.8%) |
| Prevention | Dental Visit | 2 | 74.3% (71.2% - 77.3%) | 2 | 73.6% (70.3% - 76.9%) |
| Prevention | Core preventive services for older women | 14 | 42.2% (35.1% - 49.6%) | 15 | 42.8% (35.4% - 50.1%) |
| Prevention | Cholesterol Screening | 60 | 82.2% (78.7% - 85.8%) | 16 | 80.1% (76.2% - 84.0%) |
| Prevention | Core preventive services for older men | 30 | 44.5% (36.3% - 52.8%) | 31 | 44.9% (36.8% - 53.0%) |
| Prevention | Health Insurance | 9 | 6.6% (4.8% - 8.7%) | 8 | 6.9% (5.1% - 9.1%) |
| Prevention | Mammography | 4 | 77.5% (73.6% - 81.4%) | 4 | 77.3% (73.1% - 81.3%) |

| Category | Health Domain | Crude Rate Rank | Crude Rate (95% CI) | Age-Adj. Rate Rank | Age-Adj. Rate (95% CI) |
|------------|--------------------------------|--------------------|-----------------------|-----------------------|------------------------|
| Prevention | Colorectal Cancer Screening | 20 | 74.7% (71.8% - 77.5%) | 12 | 72.6% (69.3% - 75.8%) |
| Prevention | Cervical Cancer Screening | 11 | 83.8% (81.3% - 86.4%) | 6 | 84.6% (82.2% - 87.1%) |
| Prevention | Taking BP Medication | 84 | 79.4% (75.6% - 82.8%) | 82 | 58.6% (53.1% - 64.1%) |
| Prevention | Annual Checkup | 27 | 76.5% (71.3% - 81.1%) | 5 | 74.2% (68.7% - 79.2%) |
| Disability | Mobility Disability | 10 | 10.4% (8.4% - 12.6%) | 9 | 8.7% (7.1% - 10.5%) |
| Disability | Independent Living Disability | 9 | 5.8% (4.8% - 6.9%) | 8 | 5.5% (4.6% - 6.6%) |
| Disability | Self-care Disability | 7 | 2.9% (2.4% - 3.5%) | 7 | 2.6% (2.2% - 3.1%) |
| Disability | Hearing Disability | 11 | 6.7% (5.8% - 7.8%) | 8 | 5.6% (4.8% - 6.5%) |
| Disability | Cognitive Disability | 11 | 10.3% (8.4% - 12.3%) | 11 | 10.6% (8.6% - 12.8%) |
| Disability | Vision Disability | 8 | 3.5% (2.9% - 4.2%) | 6 | 3.2% (2.7% - 3.8%) |
| Disability | Any Disability | 11 | 23.3% (19.1% - 27.8%) | 9 | 21.5% (17.6% - 25.7%) |

The 95% confidence interval shown in parentheses represents the range of 95% of CDC area estimates under different model assumptions using Monte Carlo simulations.


Appendix 3: Micropolitan Counties

Marion County falls into the group of micropolitan counties. In this data, 15 counties fall into this group. The table below lists all counties in the group, sorted by the population estimate in the CDC PLACES dataset.

| County | Population Group | Population |
|--------------------|------------------|------------|
| Clinton County | Micropolitan | 46,463 |
| Cerro Gordo County | Micropolitan | 42,706 |
| Muscatine County | Micropolitan | 42,688 |
| Marshall County | Micropolitan | 39,853 |
| Des Moines County | Micropolitan | 38,491 |
| Webster County | Micropolitan | 37,147 |
| Wapello County | Micropolitan | 35,256 |
| Marion County | Micropolitan | 33,380 |
| Lee County | Micropolitan | 33,215 |
| Mahaska County | Micropolitan | 21,984 |
| Buena Vista County | Micropolitan | 20,771 |
| Carroll County | Micropolitan | 20,692 |
| Dickinson County | Micropolitan | 17,851 |
| Clay County | Micropolitan | 16,440 |
| Jefferson County | Micropolitan | 15,647 |

Metropolitan and micropolitan areas can be found in the link and the maps below.


https://www.icip.iastate.edu/maps/refmaps/cofips




Appendix 4: County Maps

Poor General Health

Fair or poor self-rated health status among adults aged >= 18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Data Source: CDC PLACES 2023 Release, BRFSS 2021

Poor Physical Health

Physical health not good for >=14 days among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Poor Mental Health

Mental health not good for >=14 days among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Binge Drinking

Binge drinking among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Current Smoking

Current smoking among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Physical Inactivity

No leisure–time physical activity among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Sleep <7 hours

Sleeping less than 7 hours among adults aged >=18 years





Data Source: CDC PLACES 2023 Release, BRFSS 2020



Arthritis


Arthritis among adults aged >=18 years






High Blood Pressure

High blood pressure among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Cancer (except skin)

Cancer (excluding skin cancer) among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Current Asthma

Current asthma among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Coronary Heart Disease

Coronary heart disease among adults aged >=18 years





Data Source: CDC PLACES 2023 Release, BRFSS 2021



COPD


Chronic obstructive pulmonary disease among adults aged >=18 years






Depression

Depression among adults aged >=18 years





Data Source: CDC PLACES 2023 Release, BRFSS 2021



Diabetes


Diagnosed diabetes among adults aged >=18 years






High Cholesterol

High cholesterol among adults aged >=18 years who have been screened in the past 5 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Chronic Kidney Disease


Chronic kidney disease among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021





Obesity
Obesity among adults aged >=18 years







Stroke
Stroke among adults aged >=18 years






All Teeth Lost
All teeth lost among adults aged >=65 years






Lack of Health Insurance

Current lack of health insurance among adults aged 18-64 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Taking BP Medication

Taking medicine for high blood pressure control among adults aged >=18 years with high blood pressure




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Cervical Cancer Screening

Cervical cancer screening among adult women aged 21-65 years




Data Source: CDC PLACES 2023 Release, BRFSS 2020




Annual Checkup

Visits to doctor for routine checkup within the past year among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Cholesterol Screening

Cholesterol screening among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Colorectal Cancer Screening

Colorectal cancer screening among adults aged 50-75 years




Data Source: CDC PLACES 2023 Release, BRFSS 2020




Core preventive services for older men

Older adult men aged >=65 years who are up to date on a core set of clinical preventive services: Flu shot past year, PPV shot ever, Colorectal cancer screening




Data Source: CDC PLACES 2023 Release, BRFSS 2020




Core preventive services for older women

Older adult women aged >=65 years who are up to date on a core set of clinical preventive services: Flu shot past year, PPV shot ever, Colorectal cancer screening,





Data Source: CDC PLACES 2023 Release, BRFSS 2020



Dental Visit


Visits to dentist or dental clinic among adults aged >=18 years






Mammography

Mammography use among women aged 50-74 years




Data Source: CDC PLACES 2023 Release, BRFSS 2020




Hearing disability

Hearing disability among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Vision disability

Vision disability among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Cognitive disability

Cognitive disability among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Mobility disability

Mobility disability among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Self-care disability

Self-care disability among adults aged >=18 years




Data Source: CDC PLACES 2023 Release, BRFSS 2021




Independent living disability

Independent living disability among adults aged >=18 years



Data Source: CDC PLACES 2023 Release, BRFSS 2021



Any disability

Any disability among adults aged >=18 years

